



Genome Sequences of *Shewanella baltica* and *Shewanella morhuae* Strains Isolated from the Gastrointestinal Tract of Freshwater Fish

Castillo, Daniel; Gram, Lone; Dailey, Frank E.

Published in:
Genome Announcements


DOI:
[10.1128/genomeA.00541-18](https://doi.org/10.1128/genomeA.00541-18)

Publication date:
2018

Document version
Publisher's PDF, also known as Version of record

Document license:
[CC BY](#)

Citation for published version (APA):
Castillo, D., Gram, L., & Dailey, F. E. (2018). Genome Sequences of *Shewanella baltica* and *Shewanella morhuae* Strains Isolated from the Gastrointestinal Tract of Freshwater Fish. *Genome Announcements*, 6(25), 1-2. [e00541-18]. <https://doi.org/10.1128/genomeA.00541-18>



Genome Sequences of *Shewanella baltica* and *Shewanella morhuae* Strains Isolated from the Gastrointestinal Tract of Freshwater Fish

Daniel Castillo,^a Lone Gram,^b Frank E. Dailey^c

^aMarine Biological Section, University of Copenhagen, Helsingør, Denmark

^bDepartment of Biotechnology and Biomedicine, Technical University of Denmark, Kgs Lyngby, Denmark

^cDepartment of Biology, University of Wisconsin–Sheboygan, Sheboygan, Wisconsin, USA

ABSTRACT We present here the genome sequences of *Shewanella baltica* strain CW2 and *Shewanella morhuae* strain CW7, isolated from the gastrointestinal tract of *Salvelinus namaycush* (lean lake trout) and *Coregonus clupeaformis* (whitefish), respectively. These genome sequences provide insights into the niche adaptation of these specific species in freshwater systems.

Shewanella spp. are saprophytic, Gram-negative, rod-shaped members of the order *Alteromonadales*, family *Alteromonadaceae*, within the gamma subdivision of the *Proteobacteria*. They are frequently isolated from nonhuman sources, and some species (e.g., *S. algae*) are considered pathogenic in humans (1). The genus *Shewanella* is currently composed of more than 50 species that inhabit a range of aquatic environments (2, 3). The organisms potentially play key roles in environmental processes such as bioremediation of toxic elements and heavy metals (4). Also, they may serve as microbial fuel cells (5). *Shewanella* strains have also been isolated from the gastrointestinal tracts of freshwater fish species that live in environments that seasonally approach 0°C (6, 7). Here, we report the genome sequences of *S. baltica* strain CW2 and *S. morhuae* strain CW7, which were isolated as omega-3 fatty acid producers from the intestines of the freshwater fish *Salvelinus namaycush* and *Coregonus clupeaformis*, respectively.

PacBio sequencing with 100× coverage was performed as previously described for *Shewanella* sp. strain WE21 (6). Annotation was performed with the NCBI Prokaryotic Genome Annotation Pipeline (PGAP) (8). Additionally, the genomes were analyzed with the Rapid Annotations using Subsystems Technology (RAST) server (9). Specific genomic islands were recognized by MAUVE (10), secondary metabolism was analyzed by antiSMASH (11), clustered regularly interspaced short palindromic repeat (CRISPR) arrays were determined with CRISPRFinder (12), and prophage-related sequences were identified with PHASTER (13).

The final assembly for *S. baltica* strain CW2 resulted in one complete contig of 4,944,783 bp. The assembly of CW7 resulted in two large contigs totaling 4,303,233 bp. The assembly program may have failed to close the contigs due to a large repeat region at the ends. The GC contents of the genomes were 46.4 and 44%, respectively. Genome annotation for *S. baltica* resulted in 4,275 coding sequences (CDS) and 144 RNAs. *S. morhuae* had 3,764 CDSs and 137 RNAs.

In addition, both genomic sequences contained a type I secretion system (*lapBCDE*, *lapL*, *lapP*, and *rtx*); however, *S. baltica* strain CW2 had only one agglutination *rtx* toxin gene, and *S. morhuae* strain CW7 had 19 copies. Comparison of whole genomes by MAUVE with eight marine *S. baltica* strains available in the NCBI database showed that *S. baltica* strain CW2 had five genomic islands, ranging from 13.3 to 25.9 kb, that encode

Received 15 May 2018 Accepted 15 May 2018 Published 21 June 2018

Citation Castillo D, Gram L, Dailey FE. 2018. Genome sequences of *Shewanella baltica* and *Shewanella morhuae* strains isolated from the gastrointestinal tract of freshwater fish. Genome Announc 6:e00541-18. <https://doi.org/10.1128/genomeA.00541-18>.

Copyright © 2018 Castillo et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Frank E. Dailey, frank.dailey@uwc.edu.

type I restriction-modification systems, hemoglobin-like proteins, a type II restriction-modification system, glutamate synthase, and anaerobic dimethyl sulfoxide reductase (chains A and B). Although related *S. baltica* marine isolates contain one to three plasmids (14), strain CW2 contains no plasmids or genes for a type III secretion system. Although the genes for OmpA were found in both strains, only the gene for the AquaporinZ protein was found in *S. baltica*, perhaps enabling it to grow in 0% NaCl (6). Finally, *S. morhuae* harbored a megaplasmid of 113.6 kb, which encodes a glutathione-dependent pathway of formaldehyde detoxification (*gfa*, *frmA*, *fghA*, *frmR*, and *regF*). No CRISPR arrays or prophage-like elements were found in either genomic sequence. Thus, these genomic sequences can facilitate future comprehensive comparisons and phylogenetic analyses of the niche adaptation of *Shewanella* communities.

Accession number(s). The complete genome sequence of *S. baltica* strain CW2 has been deposited at DDBJ/ENA/GenBank under the accession number [CP028355](#). The version described in this paper is the first version, CP028355.1. The whole-genome shotgun project for *S. morhuae* strain CW7 has been deposited at DDBJ/ENA/GenBank under the accession number [PYSG00000000](#). The version described in this paper is the first version, PYSG01000000.

ACKNOWLEDGMENTS

We acknowledge the Institute of Genome Sciences (IGS), University of Maryland, for performing the genomic sequencing. We also acknowledge support from the University of Wisconsin–Sheboygan (F. E. Dailey), as well as previous support from Concordia University, Wisconsin, for the genome sequencing.

This work was also supported by grant 12132390 (ProAqua project) from the Innovation Fund Denmark (L. Gram and D. Castillo).

REFERENCES

- Janda JM, Abbott SL. 2014. The genus *Shewanella*: from the briny depths below to human pathogen. *Crit Rev Microbiol* 40:293–312. <https://doi.org/10.3109/1040841X.2012.726209>.
- Fonnesbech Vogel B, Venkateswaran K, Satomi M, Gram L. 2005. Identification of *Shewanella baltica* as the most important H₂S-producing species during iced storage of Danish marine fish. *Appl Environ Microbiol* 71:6689–6697. <https://doi.org/10.1128/AEM.71.11.6689-6697.2005>.
- Satomi M, Vogel BF, Gram L, Venkateswaran K. 2006. *Shewanella hafnensis* sp. nov. and *Shewanella morhuae* sp. nov., isolated from marine fish of the Baltic Sea. *Int J Syst Evol Microbiol* 56:243–249. <https://doi.org/10.1099/ijs.0.63931-0>.
- Fredrickson JK, Romine MF, Beliaev AS, Auchtung JM, Driscoll ME, Gardner TS, Nealson KH, Osterman AL, Pinchuk G, Reed JL, Rodionov DA, Rodrigues JLM, Saffarini DA, Serres MH, Spormann AM, Zhulin IB, Tiedje JM. 2008. Towards environmental systems biology of *Shewanella*. *Nat Rev Microbiol* 6:592–603. <https://doi.org/10.1038/nrmicro1947>.
- Qian F, Morse DE. 2011. Miniaturizing microbial fuel cells. *Trends Biotechnol* 29:62–69. <https://doi.org/10.1016/j.tibtech.2010.10.003>.
- Dailey FE, McGraw JE, Jensen BJ, Bishop SS, Lokken JP, Dorff KJ, Ripley MP, Munro JB. 2016. The microbiota of freshwater fish and freshwater niches contain omega-3 fatty acid-producing *Shewanella* species. *Appl Environ Microbiol* 82:218–231. <https://doi.org/10.1128/AEM.02266-15>.
- Castillo D, Gram L, Dailey FE. 2018. Complete genome sequence of *Shewanella* sp. WE21, a rare isolate with multiple novel large genomic islands. *Genome Announc* 6(16):e00277-18. <https://doi.org/10.1128/genomeA.00277-18>.
- Tatusova T, DiCuccio M, Badretdin A, Chetvernin V, Nawrocki EP, Zaslavsky L, Lomsadze A, Pruitt KD, Borodovsky M, Ostell J. 2016. Prokaryotic Genome Annotation Pipeline. *Nucleic Acids Res* 44:6614–6624. <https://doi.org/10.1093/nar/gkw569>.
- Overbeek R, Olson R, Pusch GD, Olsen GJ, Davis JJ, Disz T, Edwards RA, Gerdes S, Parrello B, Shukla M, Vonstein V, Wattam AR, Xia F, Stevens R. 2014. The SEED and the Rapid Annotation of microbial genomes using Subsystems Technology (RAST). *Nucleic Acids Res* 42:D206–D214. <https://doi.org/10.1093/nar/gkt1226>.
- Darling AE, Mau B, Perna NT. 2010. progressiveMauve: multiple genome alignment with gene gain, loss and rearrangement. *PLoS One* 5:e11147. <https://doi.org/10.1371/journal.pone.0011147>.
- Weber T, Blin K, Duddela S, Krug D, Kim HU, Brucoleri R, Lee SY, Fischbach MA, Müller R, Wohlleben W, Breitling R, Takano E, Medema MH. 2015. antiSMASH 3.0—a comprehensive resource for the genome mining of biosynthetic gene clusters. *Nucleic Acids Res* 43:W237–W243. <https://doi.org/10.1093/nar/gkv437>.
- Grissa I, Vergnaud G, Pourcel C. 2007. CRISPRFinder: a Web tool to identify clustered regularly interspaced short palindromic repeats. *Nucleic Acids Res* 35:W52–W57. <https://doi.org/10.1093/nar/gkm360>.
- Arndt D, Grant J, Marcu A, Sajed T, Pon A, Liang Y, Wishart DS. 2016. PHASTER: a better, faster version of the PHAST phage search tool. *Nucleic Acids Res* 44:W16–W21. <https://doi.org/10.1093/nar/gkw387>.
- Caro-Quintero A, Auchtung A, Deng J, Brettar I, Höfle M, Tiedje JM, Konstantinidis KT. 2012. Genome sequencing of five *Shewanella baltica* strains recovered from the oxic-anoxic interface of the Baltic Sea. *J Bacteriol* 194:1236. <https://doi.org/10.1128/JB.06468-11>.